



SEIZURE/CONVULSION - PEDIATRIC

**ALS STANDING ORDERS:**

1. For on-going seizure activity:

▶ *Turn to side, protect airway and suction when possible*

If over 12-months of age, give:

▶ *Midazolam 0.1 mg/kg IN /IM (maximum 5 mg per dose); repeat 0.1 mg/kg IN/IM one time for continued seizure activity 3 minutes after initial dose or for continued or recurrent seizure.*

If under 12-months of age, make base contact for orders and destination.

2. Pulse oximetry, if room air oxygen saturation less than 95%:

▶ *Provide high-flow Oxygen by mask or nasal cannula as tolerated.*

3. Obtain blood glucose and document finding, if blood glucose equal to or less than 60, administer one of:

▶ *10% Dextrose 5 mL/kg IV (maximum dose 200 mL).*

▶ *Glucagon 0.5 mg IM if unable to establish IV.*

*Note: IO access may be used for dextrose administration when patient is unconscious with blood glucose less than 60, unable to establish IV and there is no response to glucagon.*

4. ALS escort to nearest appropriate ERC or contact Base Hospital as needed.

Approved:

Review Dates: 05/16, 11/16, 06/17  
Final Date for Implementation: 10/01/2017  
OCEMS copyright © 2017